

Cyclization of substituted 2-(2-fluorophenylazo)azines to azino[1,2-*c*]benzo[*d*][1,2,4]triazinium derivatives

Aleksandra Jankowiak¹, Emilia Obijalska^{1,§} and Piotr Kaszynski^{*1,2}

Full Research Paper

Open Access

Address:

¹Organic Materials Research Group, Department of Chemistry, Vanderbilt University, Nashville, TN 37235, USA and ²Faculty of Chemistry, University of Łódź, Tamka 12, 91403 Łódź, Poland

Email:

Piotr Kaszynski* - piotr.kaszynski@vanderbilt.edu

* Corresponding author

§ A visiting student from the laboratory of Professor Grzegorz Mlostoń, University of Łódź, Poland.

Keywords:

DFT; heterocycles; mechanism; synthesis

Beilstein J. Org. Chem. **2013**, *9*, 1873–1880.

doi:10.3762/bjoc.9.219

Received: 24 June 2013

Accepted: 19 August 2013

Published: 16 September 2013

Associate Editor: T. J. J. Müller

© 2013 Jankowiak et al; licensee Beilstein-Institut.

License and terms: see end of document.

Abstract

Light-induced cyclization of several substituted 2-(2-fluorophenylazo)azines in the presence of Ca²⁺ ions to the corresponding triazinium derivatives is investigated experimentally and computationally. The azo derivatives of 4-methylpyridine **4** undergo facile cyclization to the corresponding triazinium **1**, and the rate of cyclization increases with increasing number of fluorine atoms at the benzene ring. No triazinium ions were obtained from azo derivatives of 4-cyanopyridine, pyrazine and pyrimidine, presumably due to their instability under the reaction conditions. The experimental results and mechanism are discussed with the aid of DFT computational results.

Introduction

Derivatives of benzo[*c*]quinolizinium [1-4] (**I**, X = Y = CH, Figure 1) and its di-aza analogue pyrido[2,1-*c*]benzo[*d*][1,2,4]triazinium (**I**, X = Y = N) belong to a class of tricyclic cations featuring general structure **I** and are of general interest because of their biological activity. For instance, it was demonstrated that the parent cation **1a** intercalates into the minor groove of DNA [5] and to be cytotoxic to cisplatin-resistant tumor cell lines [6]. Several of its derivatives containing Cl, Br, Me and OMe substituents at the benzene ring were prepared by acid-catalyzed photoinduced cyclization of phenylazopyridines [5]. Recently, we demonstrated an efficient alternative method

for the formation of **1a** and other parent tricyclic azinium cations of general structure **I** (A = B = H) by light-induced intramolecular cyclization of fluorides **II** (Figure 1) in the presence of Ca²⁺ ions [7].





Figure 1: Structures of cations **I** and precursors **II**.



A further progress in the investigation of cation **1a** and its applications as a pharmacophore or a component of organic materials requires access to functionalized derivatives, in which substituents control the properties and allow the incorporation into more complex molecular structures. As the first step towards these goals we tested the suitability of our previously described method [7] for the synthesis of several derivatives of **1a** and also for two other azines. Initially we selected CH₃, CN and F substituents to investigate the reactivity of the azines, and CH₂OH to provide a potential synthetic handle for the incorporation into more complex structures. Here we present the synthesis of four substituted derivatives of **1a** and investigate the preparation of cations derived from pyrazine **2** and pyrimidine **3** shown in Figure 2. Our experimental results are augmented with DFT computational results.



Results and Discussion

Cyclization and cation formation

Salts **1** were prepared by photocyclization of the corresponding azo derivatives **4** in the presence of Ca²⁺ ions in aqueous MeCN (Scheme 1). Diazenes **4** show weak absorption bands in the visible (e.g. 452 nm, log ε = 2.6 for **4c**), while salts **1** exhibit only a weak tailing absorption above 400 nm, as shown for **4c** and **1c** in Figure 3. Therefore irradiation was effectively conducted in Pyrex vessels.



The cyclization of 2-fluorophenylazopyridines **4b** and **4e** was conducted using a 500 W halogen lamp in the presence of Ca(OTs)₂ [7], and cations **1b** and **1e** were isolated as tosylates in 75% and 95% yield, respectively. Under these conditions diazenes **4c** and **4d** underwent cyclization, but the resulting cations **1c** and **1d** were unstable and partially decomposed during the reaction. Therefore, their cyclization was carried out with sunlight and tosylate **1c** was isolated in pure form in 90% yield. In contrast, cation **1d** partially decomposed during the workup, presumably due to the activation by means of the fluorine atoms and their electron-withdrawing effect. A pure sample of **1d** could not be isolated. A comparison of the cyclization rates of the three fluorophenylazo derivatives **4b–4d** was conducted in NMR tubes exposed to sunlight. The experiment demonstrated that the rate of cyclization increases with increasing number of fluorine atoms, and after 1 h conversion of **4b** to **1b** was 30%, **4c** to cation **1c** 80%, and **4d** to **1d** was complete. Full conversion of **4b** to **1b** was achieved after 5 h. It should be pointed out that the cyclization and formation of the cations requires both light and Ca²⁺ ions, and no product formation was observed if one of them is absent. While in most experiments Ca(OTs)₂ served as a source of Ca²⁺, in some other CaCl₂ was used, and the Cl[−] anion was later replaced by TsO[−].

Attempts to cyclize the cyano derivative **4f**, using a halogen lamp were unsuccessful and after 3 h only decomposition products were observed by ¹H NMR along with some residual starting **4f**. No characteristic NMR pattern that could be ascribed to the cation **1f** was detected. Similarly unsuccessful were the attempts to cyclize azopyrazine **5** (halogen lamp/ice bath or sunlight, Figure 4) and azopyrimidine **6** (halogen lamp/reflux); instead complex mixtures of products were obtained.




Figure 4: Structures of *trans* azo derivatives 5-*E* and 6-*E*.

Synthesis of azo precursors


Azo compounds **4b–4e** were prepared in 38%–90% yield using method A (Scheme 2) in which 2-nitroso-4-picoline (**7**) [8] was condensed with 2-fluoroaniline (**8a**) or its derivatives **8c–8e**. The 2-fluoro-5-hydroxymethylaniline (**8e**) was obtained according to a literature procedure [9]. 4-Cyano derivative **4f** was prepared in 50% yield by method B [7,10] involving condensation of 2-amino-4-cyanopyridine with 1-fluoro-2-nitrosobenzene (**9**) [7]. Azoazines **5** and **6** were obtained in a similar way by condensation of **9** with 2-aminopyrazine and 2-aminopyrimidine, respectively, and their preparation was described before [11].

Interestingly, the trifluoro derivative **4d** had exceptionally limited thermal stability: After heating at 60 °C in CDCl₃ it was converted to another species also containing the picoline fragment, according to the ¹H NMR analysis. The new compound did not undergo photocyclization and was not analyzed further.

Mechanistic and computational analyses


The proposed mechanism for the formation of the triazinium cations from fluorophenylazoazines involves a photo-induced *trans*-to-*cis* isomerization of the azo group followed by Ca²⁺-assisted cyclization (Scheme 3) [7], which, in principle, can proceed either thermally or photochemically. Thus, the success of the reaction depends on i) the efficient access to the *Z*-isomer, and ii) the relative rates of cyclization vs back

isomerization to the *E*-isomer. The supply of the requisite *Z*-isomers of the azoazines is best accomplished by photoisomerization of the *E*-isomers, since the *Z*-isomers cannot be populated thermally due to the high *E* → *Z* thermal isomerization barrier ($\Delta G_{298}^\ddagger > 35$ kcal/mol) [7]. The *Z*-forms however, thermally and also photochemically isomerize back to the more stable *E*-isomers, and the rates are substituent depended [12]. Thus, with the broadband irradiation a photostationary state establishes providing a constant ratio of *Z*- to *E*-isomers.



Scheme 3: Formation of cations **1** from diazenes **4**.

Previous analyses suggested [7] that the cyclization of **4-Z** involves 6 π electrons leading to non-aromatic product **10**, which upon Ca²⁺-assisted departure of the fluoride undergoes aromatization and formation of cation **1**. In this work, the theoretical analysis is concentrated on the effect of substituents and on the type of azine on the formation of the cations. Therefore the cyclization of *cis*-azoazines **4-Z–6-Z** was investigated at B3LYP/6-311+G(2d,p)//B3LYP/6-311G(2d,p) level of theory in MeCN as dielectric medium without participation of metal ions. The computational studies were expanded by inclusion of **4g-Z–4i-Z** for a better understanding of the structure–reactivity



Scheme 2: Synthesis of azo precursors. Reagents and conditions: i) AcOH cat, CH₂Cl₂, rt, 25 h; ii) toluene, 50% NaOH, 50 °C, 25 min; iii) NCS, Me₂S, CH₂Cl₂, –20 °C; iv) mCPBA, CH₂Cl₂, 0 °C; v) Oxone®, CH₂Cl₂/H₂O rt, 25 h.

relationship. The resulting free energy differences between the Z-isomers, relevant to the cyclization process, and the cyclic products, and free energy of activation for a series of diazenes are listed in Table 1 and Table 2.

Table 1: Calculated free energy activation and change (kcal/mol) for the cyclization of *cis*-azopyridine **4-Z** to the nonaromatic product **10** in MeCN (Scheme 3).^a

	R	X	Y	Z	ΔG_{298}^\ddagger ^b	ΔG_{298}^c
a	H	H	H	H	22.4	5.7
b	Me	H	H	H	23.1	5.7
c	Me	F	H	H	18.5	2.4
d	Me	F	F	H	18.9	4.6
f	CN	H	H	H	23.3	5.2
g	OMe	H	H	H	23.5	4.5
h	Me	H	H	OMe	23.2	3.9
i	Me	H	H	CN	19.0	4.8

^aCalculations at B3LYP/6-311+G(2d,p)//B3LYP/6-311G(2d,p) level with MeCN as dielectric medium. ^bEnergy of the transition state **10-TS**. ^cEnergy difference between product **10** and azopyridine **4-Z**.

Table 2: Calculated free energy activation and change (kcal/mol) for cyclization of *cis*-azoazines to the nonaromatic product in MeCN.^a

reaction	ΔG_{298}^\ddagger ^b	ΔG_{298}^c
4a-Z → 10a	22.4	5.7
5-Z → 11	18.7	2.7
6-Z → 12	25.0	7.5

^aCalculations at B3LYP/6-311+G(2d,p)//B3LYP/6-311G(2d,p) level in MeCN as dielectric medium. ^bEnergy of the transition state. ^cEnergy difference between the cyclic product and azoazine.

The results in Table 1 and Table 2 show that the free energy of activation, ΔG_{298}^\ddagger , required for the cyclization of all considered azoazines is in a range of 18–25 kcal/mol, and the process

is endergonic by about 5 kcal/mol. Thus, for all investigated compounds the cyclization is thermally accessible and also thermally reversible at ambient temperature. These computational results are consistent with experimental observations, that all azoazines reacted, however only 5 cations (**1a–1e**) were isolated. The remaining cations, **1f**, **2** and **3**, could have been formed, but were presumably unstable under the reaction conditions, as already observed for **1d**. Also consistent with the computational results is the observation that the formation of **1** requires a fluoride scavenger, apparently for the aromatization of the cyclic product.

The analysis of species involved in the cyclization demonstrates that the C···N and C–F distances change similarly for all azoazines in the process regardless of the substituent. Thus, the C···N distance decreases by 1.232 Å from 3.150 ± 0.03 Å in the *cis*-azoazine to 1.918 ± 0.009 Å in the transition state (TS) and then to 1.457 ± 0.004 Å in the cycloadduct, as shown for the cyclization of **4c** to **10c** in Figure 5. At the same time the C–F distance increases only by about 2% from 1.350 ± 0.002 Å in the *cis*-azoazine to 1.374 ± 0.003 Å in the TS. A more significant increase in the C–F distance, by 7%, is observed in the cyclic product (1.445 ± 0.005 Å).

A detailed analysis of the DFT results for the azopyridine series demonstrates that the rate of cyclization only correlates with the character of the substituent at the benzene ring and does not depend on the pyridine ring. For a simple nucleophilic aromatic *ipso*-substitution reaction (NAS) electron-donating groups in the C(4) position of the pyridine ring in **4** should increase the reaction rate. Computational results demonstrate the contrary and as the electron donation of the substituent R increases in the order H (**4a**) < CH₃ (**4b**) < OCH₃ (**4g**) so does the activation energy (Table 1). In contrast, increasing electron withdrawing ability of the substituent Z in the *para*-position to the




Figure 5: B3LYP/6-311G(2d,p)-optimized geometries for structures involved in cyclization of **4c** to **1c**.

leaving fluorine atom increases the reaction rate as observed in a series **4h** ($Z = \text{OCH}_3$, $\Delta G^\ddagger_{298} = 23.2$ kcal/mol), **4a** ($Z = \text{H}$, $\Delta G^\ddagger_{298} = 22.4$ kcal/mol), and **4i** ($Z = \text{CN}$, $\Delta G^\ddagger_{298} = 19.0$ kcal/mol) in Table 1.

For further probing of the cyclization mechanism, the mildly activating [13,14] N=N-bridging group in **4c-Z** was replaced by the non-activating CH=CH group in **13-Z** (Figure 6). A computational analysis demonstrated that this structural modification resulted in a modest increase of activation energy ΔG^\ddagger_{298} for the cyclization by 3.1 kcal/mol. However, replacement of the bridging π -bond with the aliphatic CH_2CH_2 group in compound **14** prevents the cyclization process, and the analogous cyclic product could not be located on the potential energy surface. These results indicate that the cyclization process involves 6 π electrons, which allows the formation of the non-zwitterionic product such as **10**. Comparison of cyclization of **4c-Z** and **13-Z** further demonstrates that the activation energy is lower for the azene than for the analogues stilbene, which presumably is related to the electronegativity of the bridging group.




Figure 6: Structures of three close analogues.

The analysis of the data of three fluoro derivatives **4b–4d** in Table 1 demonstrates that the addition of one or two fluorine atoms at the benzene ring in **4b** significantly lowers the activation barrier for the cyclization, by about 4.5 kcal/mol, for **4c-Z** and **4d-Z**, which is consistent with the accelerated formation of cations **1c** and **1d** observed experimentally. Interestingly, a bigger effect is predicted for the 2,4-difluorophenyl derivative **4c** than for the 2,4,6-trifluorophenyl analogue **4d**.

Among the three parent azoazines the least unfavorable cyclization is calculated for the pyrazine derivative **5-Z** ($\Delta G_{298} = 2.7$ kcal/mol), which also has the lowest TS energy ($\Delta G^\ddagger_{298} = 18.7$ kcal/mol, Table 2). On the other hand, the cyclization of pyrimidine **6-Z** is most endergonic and has the highest TS energy among all azoazines considered in this work. These significant differences between the three parent azoazines can be explained with lower aromaticity of the pyrazine ring [15], and with the electropositive character of the N(1) position

of the pyrimidine ring, when compared to the pyridine analogue. These two factors may also contribute to the higher reactivity of cations **2** and **3** and hence their apparent instability under the reaction conditions.

Conclusion

A DFT computational analysis indicates that the cyclization of *cis*-fluorophenylazoazines can occur thermally at room temperature for all considered compounds. The reaction is endergonic with 2.4–7.5 kcal/mol, and the formation of the triazinium cations is accomplished by aromatization of the cyclic product by Ca^{2+} ion-assisted removal of the fluoride. These results are supported by experimental data, which demonstrate a facile conversion of all azoazines under the reaction conditions, although only 5 out of 8 triazinium salts were isolated. The remaining three, all of them containing a CN substituent and derived from pyrazine and pyrimidine, are presumably too reactive under the reaction conditions to be isolated in pure form.

A brief computational investigation of the mechanism demonstrates the importance of the unsaturated bridge in the cyclization step and points towards a 6π -electrocyclization rather than to a simple nucleophilic aromatic displacement mechanism.

Among the triazinium salts prepared in this work is the hydroxymethyl derivative **1e**, which, in principle, can be incorporated into more complex molecular structures.

Computational Details

The quantum-mechanical calculations were carried out at B3LYP/6-311G(2d,p) level of theory using the Gaussian 09 suite of programs [16]. The geometry optimizations were undertaken using tight convergence limits and with no symmetry constraints. Vibrational frequencies were used to characterize the nature of the stationary points. Zero-point energy (ZPE) corrections were scaled by 0.9806 [17]. Transition state structures for cyclization reactions were located using the STQN method [18] requested with the QST2 keyword and default convergence criteria. Final energies for each optimized structure were calculated with the B3LYP/6-311+G(2d,p)//B3LYP/6-311G(2d,p) method in MeCN as dielectric medium using the IPCM model [19] requested with the SCRF=IPCM keyword (epsilon = 36.64).

Experimental

General remarks

Melting points are uncorrected. NMR spectra were recorded at 300 MHz (^1H) and 100 MHz (^{13}C) in CD_3CN , $\text{DMSO}-d_6$ or CDCl_3 and chemical shifts are referred to the solvent peaks (1.96, 2.54 and 7.26 ppm, respectively).

Photocyclization and preparation of salts 1.

General Procedure

Method A [7]: Azo compound **4** (1.0 mmol) and calcium *p*-toluenesulfonate (190 mg 0.5 mmol) or calcium chloride (56 mg, 0.5 mmol) were dissolved in a mixture of MeCN/H₂O (9:1, 30 mL). The resulted solution was irradiated with a 500 W halogen lamp and gently refluxed until TLC control showed full conversion of the substrate (about 1.5 h). The solvents were evaporated. The residue was dried in a desiccator over P₂O₅ (12 h). The solid was washed with CH₂Cl₂, the residue was dissolved in hot MeCN and filtered. The solvent was evaporated and the crude product was recrystallized from aqueous MeCN.

Method B: The reaction mixture prepared as in method A was exposed to intense sunlight, while stirring until TLC control showed full conversion of the substrate (about 1.5 h). The work-up of the reaction mixture was as described in method A.

3-Methylpyrido[2,1-*c*][1,2,4]benzotriazin-11-ium *p*-toluenesulfonate (1b): Method A, yield 95%: ¹H NMR (300 MHz, CD₃CN) δ 2.33 (s, 3H), 2.98 (s, 3H), 7.16 (d, *J* = 7.8 Hz, 2H), 7.60 (d, *J* = 8.0 Hz, 2H), 8.35 (t, *J* = 7.8 Hz, 1H), 8.43 (d, *J* = 7.5 Hz, 1H), 8.49 (t, *J* = 8.1 Hz, 1H), 8.83 (d, *J* = 8.8 Hz, 1H), 9.03 (d, *J* = 8.0 Hz, 1H), 9.12 (s, 1H), 9.87 (d, *J* = 7.0 Hz, 1H); HRMS *m/z* calcd for C₁₂H₁₀N₃, 196.0869; found, 196.0892.

9-Fluoro-3-methylpyrido[2,1-*c*][1,2,4]benzotriazin-11-ium *p*-toluenesulfonate (1c): Method B, yield 90%: mp >190 °C dec.; ¹H NMR (300 MHz, CD₃CN) δ 2.35 (s, 3H), 2.96 (s, 3H), 7.19 (d, *J* = 7.5 Hz, 2H), 7.60 (d, *J* = 7.1 Hz, 2H), 8.11 (t, *J* = 7.8 Hz, 1H), 8.44 (d, *J* = 6.7 Hz, 1H), 8.64 (dd, *J*₁ = 9.2 Hz, *J*₂ = 2.2 Hz, 1H), 9.00–9.60 (m, 1H), 9.11 (s, 1H), 9.73 (d, *J* = 6.8 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 20.8, 21.8, 104.1 (d, *J* = 30 Hz), 121.8 (d, *J* = 25 Hz), 125.5, 128.0, 129.7, 130.2, 132.3, 135.7 (d, *J* = 11 Hz), 137.5, 138.9, 141.9, 145.8, 160.6; UV–vis (MeCN) λ_{max} (log ε) 225 (4.20), 254 (4.41), 262 sh (4.38), 365 (3.98), 379 sh (3.81); anal. calcd for C₁₉H₁₆FN₃O₃S: C, 59.21; H, 4.18; N, 10.90; found: C, 58.92; H, 4.09; N, 10.95.

7,9-Difluoro-3-methylpyrido[2,1-*c*][1,2,4]benzotriazin-11-ium chloride (1d): Method B, yield 40–80% (based on NMR): ¹H NMR (300 MHz, CD₃CN) δ 2.98 (s, 3H), 7.99 (t, *J* = 8.1 Hz, 1H), 8.49 (d, *J* = 5.5 Hz, 1H), 8.56 (d, *J* = 8.9 Hz, 1H), 9.16 (s, 1H), 9.77 (d, *J* = 6.8 Hz, 1H).

8-Hydroxymethyl-3-methylpyrido[2,1-*c*][1,2,4]benzotriazin-11-ium *p*-toluenesulfonate (1e): Method A, yield 75%: mp >150 °C dec.; ¹H NMR (300 MHz, DMSO-*d*₆) δ 2.32 (s,

3H), 2.98 (s, 3H), 4.97 (d, *J* = 5.0 Hz, 2H), 5.92 (t, *J* = 5.5 Hz, 1H), 7.14 (d, *J* = 7.8 Hz, 2H), 7.50 (d, *J* = 8.0 Hz, 2H), 8.49 (d, *J* = 8.5 Hz, 1H), 8.64 (d, *J* = 5.5 Hz, 1H), 8.95 (s, 1H), 9.17 (d, *J* = 8.9 Hz, 1H), 9.35 (s, 1H), 10.36 (d, *J* = 6.8 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 20.8, 21.6, 61.6, 116.4, 121.8, 125.5, 128.0, 128.1, 129.6, 130.2, 131.9, 136.6, 137.6, 141.2, 142.0, 145.7, 148.2, 159.2; anal. calcd for C₂₀H₁₉N₃O₄S: C, 60.44; H, 4.82; N, 10.57; found: C, 60.15; H, 4.78; N, 10.47.

Azo compounds 4. A general procedure

Method A. To the solution of amine (1.0 mmol) in dry CH₂Cl₂ (2 mL), 2-nitroso-4-picoline (**7**, 1.0 mmol) was added followed by a catalytic amount (1 drop) of acetic acid. The reaction mixture was stirred at rt for 24 h, protected from a light. The solvent was evaporated and the residue was purified on a silica gel plug (CH₂Cl₂/EtOAc, 5:1) to give the corresponding azo compound as orange-red crystals. Analytically pure samples were obtained by recrystallization (hexane/CH₂Cl₂).

2-(2-Fluorophenylazo)picoline (4b): Yield: 83%: mp 78–80 °C; ¹H NMR (300 MHz, CDCl₃) δ 2.48 (s, 3H), 7.18–7.34 (m, 3H), 7.48–7.56 (m, 1H), 7.64 (s, 1H), 7.90 (t, *J* = 7.7 Hz, 1H), 8.61 (d, *J* = 4.4 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 21.2, 114.5, 117.1 (d, *J* = 20 Hz), 117.9, 124.3 (d, *J* = 4 Hz), 126.5, 133.7 (d, *J* = 4 Hz), 140.4 (d, *J* = 6 Hz), 149.3, 149.8, 160.7 (d, *J* = 258 Hz), 163.3; anal. calcd for C₁₂H₁₀FN₃: C, 66.97; H, 4.68; N, 19.52; found: C, 66.93; H, 4.66; N, 19.41.

2-(2,4-Difluorophenylazo)picoline (4c): Yield 90%: mp 99–101 °C; ¹H NMR (300 MHz, CD₃CN) δ 2.48 (s, 3H), 7.14 (t, *J* = 8.8 Hz, 1H), 7.26 (ddd, *J*₁ = 11.3 Hz, *J*₂ = 8.7 Hz, *J*₃ = 2.6 Hz, 1H), 7.37 (d, *J* = 5.0 Hz, 1H), 7.60 (s, 1H), 7.91 (dd, *J*₁ = 15.3 Hz, *J*₂ = 8.8 Hz, 1H), 8.58 (d, *J* = 4.8 Hz, 1H); ¹H NMR (300 MHz, CDCl₃) δ 2.48 (s, 3H), 6.92–7.08 (m, 2H), 7.26 (d, 1H), 7.63 (s, 1H), 7.94 (d, *J* = 8.0 Hz, 1H), 7.99 (d, *J* = 8.6 Hz, 1H), 8.60 (d, *J* = 4.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 21.1, 105.1 (t, *J* = 24 Hz), 112.0 (dd, *J*₁ = 27 Hz, *J*₂ = 4 Hz), 114.5, 119.3 (d, *J* = 10 Hz), 126.5, 137.3 (dd, *J*₁ = 7 Hz, *J*₂ = 4 Hz), 149.2, 149.8, 161.2 (dd, *J*₁ = 260 Hz, *J*₂ = 7 Hz), 163.1, 165.3 (dd, *J*₁ = 255 Hz, *J*₂ = 12 Hz); UV–vis (MeCN) λ_{max} (log ε) 225 (4.02), 322 (4.22), 541 (6.61); anal. calcd for C₁₂H₉F₂N₃: C, 61.80; H, 3.89; N, 18.02; found: C, 61.79; H, 3.79; N, 17.92.

2-(2,4,6-Trifluorophenylazo)picoline (4d): Yield 38%: ¹H NMR (300 MHz, CD₃CN) δ 2.49 (s, 3H), 6.85 (t, *J* = 8.8 Hz, 2H), 7.26 (d, 1H), 7.56 (s, 1H), 8.58 (d, *J* = 4.9 Hz, 1H); ¹H NMR (300 MHz, CDCl₃) δ 2.48 (s, 3H), 7.09 (t, *J* = 9.3 Hz, 2H), 7.26 (d, 1H), 7.61 (s, 1H), 8.61 (d, *J* = 4.8 Hz,

1H); HRMS m/z calcd for $C_{12}H_9F_3N_2$, 252.0749; found, 252.0738.

2-(2-Fluoro-5-hydroxymethylphenylazo)picoline (4e): Yield 69%; mp 154–155 °C; 1H NMR (300 MHz, $CDCl_3$) δ 2.48 (s, 3H), 4.73 (s, 2H), 7.28–7.32 (m, 1H), 7.52–7.60 (m, 1H), 7.64 (s, 1H), 7.88 (d, $J = 7.1$ Hz, 1H), 7.60 (d, $J = 4.9$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 21.2, 64.0, 114.3, 115.9, 117.2 (d, $J = 20$ Hz), 126.6, 132.2 (d, $J = 8$ Hz), 137.6 (d, $J = 4$ Hz), 140.0 (d, $J = 7$ Hz), 149.1, 150.0, 160.0 (d, $J = 257$ Hz), 163.1; anal. calcd for $C_{13}H_{12}FN_3O$: C, 63.66; H, 4.93; N, 17.13; found: C, 63.94; H, 4.88; N, 17.20.

4-Cyano-2-(2-fluorophenylazo)pyridine (4f): Method B. 4-Cyano-2-aminopyridine (238 mg, 2.0 mmol) was dissolved in toluene (3 mL) and a 50% aqueous solution of NaOH (1.5 mL) and 2-fluoronitrosobenzene [7] (**9**, 293 mg, 2.3 mmol) was added. The mixture was vigorously stirred at 50 °C for 25 min. After cooling, water was added, and the mixture was extracted with CH_2Cl_2 . Extracts were dried over Na_2SO_4 , and the solvent was evaporated. The residue was purified by column chromatography (SiO_2 , hexane/ CH_2Cl_2) to give 230 mg (50% yield) of azo derivative **4f** as an orange solid: 1H NMR (300 MHz, CD_3CN) δ 7.34 (t, $J = 7.8$ Hz, 1H), 7.41 (t, $J = 9.7$ Hz, 1H), 7.62–7.72 (m, 1H), 7.76–7.88 (m, 2H), 8.02 (s, 1H), 8.89 (d, $J = 4.9$ Hz, 1H); 1H NMR (300 MHz, $CDCl_3$) δ 7.27–7.41 (m, 2H), 7.56–7.61 (m, 1H), 7.66 (d, $J = 5.0$ Hz, 1H), 8.05 (s, 1H), 8.93 (d, $J = 4.9$ Hz, 1H); ^{13}C NMR (100 MHz, CD_3CN) δ , 116.8, 117.0, 118.2, 118.3, 123.2, 125.6 (d, $J = 6$ Hz), 127.9, 135.9 (d, $J = 8$ Hz), 140.7 (d, $J = 7$ Hz), 151.4, 161.3 (d, $J = 257$ Hz), 164.0; anal. calcd for $C_{12}H_7FN_4$: C, 63.71; H, 3.12; N, 24.77; found: C, 63.81; H, 3.14; N, 24.61.

Supporting Information

Supporting Information File 1

General methods and synthetic procedures.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-9-219-S1.pdf>]

Acknowledgements

Financial support for this work was received from the Petroleum Research Funds (PRF-47243-AC4) and the National Science Foundation (OISE-0532040 and CHE-1214104).

References

- Norez, C.; Bilan, F.; Kitzis, A.; Mettey, Y.; Becq, F. *J. Pharmacol. Exp. Ther.* **2008**, *325*, 89–99. doi:10.1124/jpet.107.134502
- Roomans, G. M. *Am. J. Respir. Med.* **2003**, *2*, 413–431. doi:10.1007/BF03256668
- Prost, A.-L.; Dérand, R.; Gros, L.; Becq, F.; Vivaudou, M. *Biochem. Pharmacol.* **2003**, *66*, 425–430. doi:10.1016/S0006-2952(03)00289-2
- Galiotta, L. J. V.; Springsteel, M. F.; Eda, M.; Niedzinski, E. J.; By, K.; Haddadin, M. J.; Kurth, M. J.; Nantz, M. H.; Verkman, A. S. *J. Biol. Chem.* **2001**, *276*, 19723–19728. doi:10.1074/jbc.M101892200
- Sinan, M.; Panda, M.; Ghosh, A.; Dhara, K.; Fanwick, P. E.; Chattopadhyay, D. J.; Goswami, S. *J. Am. Chem. Soc.* **2008**, *130*, 5185–5193. doi:10.1021/ja710211u
- Garza-Ortiz, A.; den Dulk, H.; Brouwer, J.; Kooijman, H.; Spek, A. L.; Reedijk, J. *J. Inorg. Biochem.* **2007**, *101*, 1922–1930. doi:10.1016/j.jinorgbio.2007.05.008
- Jankowiak, A.; Objalska, E.; Kaszynski, P.; Pieczonka, A.; Young, V. G., Jr. *Tetrahedron* **2011**, *67*, 3317–3327. doi:10.1016/j.tet.2011.03.023
- Taylor, E. C.; Tseng, C. P.; Rampal, J. B. *J. Org. Chem.* **1982**, *47*, 552–555. doi:10.1021/jo00342a035
- Carroll, W. A.; Altenbach, R. J.; Bai, H.; Brioni, J. D.; Brune, M. E.; Buckner, S. A.; Cassidy, C.; Chen, Y.; Coghlan, M. J.; Daza, A. V.; Drizin, I.; Fey, T. A.; Fitzgerald, M.; Gopalakrishnan, M.; Gregg, R. J.; Henry, R. F.; Holladay, M. W.; King, L. L.; Kort, M. E.; Kym, P. R.; Millicic, I.; Tang, R.; Turner, S. C.; Whiteaker, K. L.; Yi, L.; Zhang, H.; Sullivan, J. P. *J. Med. Chem.* **2004**, *47*, 3163–3179. doi:10.1021/jm030356w
- Campbell, N.; Henderson, A. W.; Taylor, D. *J. Chem. Soc.* **1953**, 1281–1285. doi:10.1039/jr9530001281
- Objalska, E.; Kaszynski, P.; Jankowiak, A.; Young, V. G. *Polyhedron* **2011**, *30*, 1339–1348. doi:10.1016/j.poly.2011.02.023
- García-Amorós, J.; Velasco, D. *Beilstein J. Org. Chem.* **2012**, *8*, 1003–1017. doi:10.3762/bjoc.8.113
- Manka, J. T.; McKenzie, V. C.; Kaszynski, P. *J. Org. Chem.* **2004**, *69*, 1967–1971. doi:10.1021/jo0302399
- Fryszkowska, A.; Tilford, R. W.; Guo, F.; Kaszynski, P. *Tetrahedron* **2005**, *61*, 2327–2333. doi:10.1016/j.tet.2005.01.023
- Katritzky, A. R.; Barczynski, P.; Musumarra, G.; Pisano, D.; Szafran, M. *J. Am. Chem. Soc.* **1989**, *111*, 7–15. doi:10.1021/ja00183a002
- Gaussian 09*, Revision A.02; Gaussian, Inc.: Wallingford, CT, 2009.
- Scott, A. P.; Radom, L. *J. Phys. Chem.* **1996**, *100*, 16502–16513. doi:10.1021/jp960976r
- Peng, C.; Ayala, P. Y.; Schlegel, H. B.; Frisch, M. J. *J. Comput. Chem.* **1996**, *17*, 49–56. doi:10.1002/(SICI)1096-987X(19960115)17:1<49::AID-JCC5>3.3.CO;2-#
- Foresman, J. B.; Keith, T. A.; Wiberg, K. B.; Snoonian, J.; Frisch, M. J. *J. Phys. Chem.* **1996**, *100*, 16098–16104. doi:10.1021/jp960488j

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
[doi:10.3762/bjoc.9.219](https://doi.org/10.3762/bjoc.9.219)