



AmericasBarometer, 2014

Technical Information

<i>Country</i>	<i>Year</i>	<i>Sample Size</i>	<i>Weighted/Unweighted</i>	<i>Fieldwork dates</i>
Suriname	2014	4,000	Weighted	June 21 st to August 25 th

LAPOP AmericasBarometer 2014 round of surveys

In its effort to collect the best quality data possible and therefore produce the highest quality studies, the Latin American Public Opinion Project (LAPOP) adopted a new sample design for the AmericasBarometer 2012 round of surveys, which was also employed in 2014. The two main reasons for this decision to change the sample design from that which was used in the 2004-2010 period were: (1) updating the sample designs to reflect the population changes as revealed by recent census information, and (2) standardizing the sample sizes at the level of the municipality in order to both reduce the variance and provide a basis for using multi-level analysis drawing on municipal data. This change in the sample design makes the sample representative by municipality type¹, to enable the use of the municipality as a unit of analysis for multilevel statistical analysis. Details of the revisions are found in the description of the 2012 AmericasBarometer surveys.

In 2013 LAPOP entered into a Memorandum of Understanding (MOU) with the Institute for Social Research at the University of Michigan for assistance in and advice on the AmericasBarometer. One of the world's leading experts in sample design methodology, Dr. Jim Lepkowski, and his graduate students, advised us throughout the process. Over the course of a

¹ The new sample design included three different strata of municipalities classified according to their size. Municipalities were grouped in sizes appropriate for the country. One common grouping was (1) Municipalities with less than 25,000 inhabitants, (2) Municipalities with between 25,000 and 100,000 inhabitants, (3) Municipalities with more than 100,000 inhabitants.

year we worked with Dr. Lepkowski and his team of graduate students to review each previously developed sample design and to secure their input and advice on new designs. Our colleagues at the University of Michigan, confirmed that LAPOP had already been following the best practices, within the limits of resources at our disposal, in its sample design. Our own review of the major update we carried out in 2012 sample design left us pleased in almost every respect. The effort to obtain a standard sample size per municipality/canton/parish did not have any adverse impact on intra-class correlation levels, yet has given us a basis for calculating context effects at the local level. In some particular cases, however, in the 2014 round we requested country teams to conduct specific alterations, like updating their sampling frame to take into consideration (if available) the new 2010-2011 national census information. We also asked teams to verify that the 2012 sample design continues to reflect and represent each country population structure and distribution.

Finally, after several rounds of consultations and technical discussions with experts at the ISR at the University of Michigan on how to update the 2012 samples for the 2014 round of surveys, LAPOP requested that countries to update their samples at the block level while retaining the same primary and sub-stratification units (i.e., *Estratopri, Municipalities and Census Segments*) that were included in the 2012 sample. This means that users of prior AmericasBarometer surveys can do so knowing that the designs across time remain very similar, if not identical. Countries that had new population census available and did not experience significant population shifts or changes in their population distribution were asked to replicate the 2012 sample using the latest census information available and to replace the sampling points at the block level.

With respect to *data collection*, in the 2014 round of the AmericasBarometer we expanded the use of handheld electronic devices. For the first time, we employed for data collection the “Adgys”[©] questionnaire app designed by our partners in Cochabamba, Bolivia. The use of electronic devices for interviews and data entry in the field reduces data entry errors, supports the use of multiple languages, and permits LAPOP to track on a daily basis the progress of the survey, down to the location of interviews (which are monitored in real time, or nearly real time, but not recorded into the public datasets in order to preserve respondents’ privacy) and the timing of the interviews. The team in Bolivia worked long hours to program the samples and questionnaires into the Adgys platform for the 20 countries in which we used this technology. In the remaining 6 countries we continued our use of PDAs and a Windows Mobile-based software application supported by our hardworking partners at the University of Costa Rica.

The remaining pages of this technical note describe the sample design of the AmericasBarometer 2014 survey in Suriname.

Suriname 2014 AmericasBarometer Round

This survey was carried out between June 21st and August 25th of 2014, as part of the LAPOP AmericasBarometer 2014 wave of surveys. The 2014 survey was conducted by Vanderbilt University with the field work being carried out by DataFruit. The 2014 AmericasBarometer received generous support from many sources, including USAID, UNDP, IADB, Vanderbilt U., Princeton U., Université Laval, U. of Notre Dame, among others.

The project used a national probability sample design of voting-age adults, with a total N of 4,000 people. It involved face-to-face interviews conducted in Dutch, Sranan Tongo and English. The survey used a complex sample design, taking into account stratification and clustering, and weighting.

The Suriname 2014 survey included an oversample conducted in the Nation's Capital City of Paramaribo. The original 2012 sample size in Paramaribo (of 520 cases) was expanded in the 2014 round to 3,005 cases, with the selection of Enumeration Districts done using PPS at the Ressorst level. The oversample was drawn using a new software developed by LAPOP, called LASSO[©] (LAPOP Survey Sample Optimizer). This proprietary software allows estimating the number of dwellings in a given region using satellite images in the public domain, and then using a probabilistic method to locate sample segments (i.e., clusters) to draw a sample.

The sample consists of five strata representing the five main geographical regions: Paramaribo, Wanica/Para, Nickerie/Coronie/Saramancca, Commewijne/Marowijne, and Brokopondo/Sipaliwini. Since some of the regions have a small population, a larger sample of respondents from these regions was drawn, and sample weights were incorporated to reflect the actual known distribution of the population between the five regions. The sample must be weighted to produce a representative national result. Each stratum was further sub-stratified by urban and rural areas. Respondents were selected in clusters of 6 interviews.

Table 1 shows the unweighted sample size in each of the five regions (strata) and by municipality size in 2012 and 2014.

Table 1: Sample sizes by Strata and Municipality Size in the 2012 and 2014 AmericasBarometer Survey in Suriname

Strata	Unweighted Sample Size 2012	Unweighted Sample Size 2014
Paramaribo	520	3,005
Wanica / Para	306	321
Nickerie / Coronie / Saramancca	240	241
Commewijne / Marowijne	215	219
Brokopondo / Sipaliwini	211	214
Total	1,492	4,000
Size of Municipality		
More than 10,000 inhabitants	-	3,149
Between 3,000 and 10,000 inhabitants	-	515
Less than 3,000 inhabitants	-	336
Total	-	4,000

The sample consists of 163 primary sampling units and 662 final sampling units. A total of 3,011 respondents were surveyed in urban areas and 989 in rural areas. The estimated margin of error for the survey is ± 1.8 .

Quotas for gender and age were adopted since multiple recalls in a national sample such as this are impractical from a cost standpoint. Our experience shows that even three recalls leave the sample with a notable gender imbalance (more women than men). Rather than have to include post-hoc weights to adjust for this sample error, we resolve the problem in the field via quotas.

Weighting of the Suriname datasets

The AmericasBarometer Suriname sample of Suriname is not self-weighted. The dataset contains a variable called WT which is the “country weight” variable. The variable “WEIGHT1500” should be activated to produce representative national results. When using this dataset for cross-country comparisons, in order to give each country in the study an identical weight in the pooled sample, LAPOP reweights each country data set in the merged files so that each country has an N of 1,500. In SPSS this is done via the “weight” command.

The questionnaire used in this round and the datasets can be found on the LAPOP website: www.AmericasBarometer.org.