

Multi-Spectral Thermal Imaging with Inversely Designed Optical Filters for Material Recognition





Thiago S. Arnaud¹, Mingze He², Josh Nordlander³, Jon-Paul Maria³, Joshua D. Caldwell²

¹Department of Mechanical Engineering, Florida International University, Miami, FL, ²Department of Mechanical Engineering, Vanderbilt University, Nashville, TN, ³Pennsylvania State University, State College, PA




Background

- All materials above absolute zero emit thermal radiation.
- Objects with temperatures near or above room temperature (~300K) have peak emission at 7 μm -14 μm wavelengths.
- A **Blackbody** is an object that emits a broadband of thermal radiation defined by Planck's Blackbody Radiation Law.
- A **graybody** is an object that emits light with the same spectral shape, but scaled to lower powers by a factor called the **emissivity**.




Motivation

Emitted Power Measurement Process



Approximating SiC's Emitted Power




- Real materials don't exactly follow Planck's Blackbody Radiation Law
- Instead, material properties such as vibrations and conductivity change the IR absorption/emission properties
- Thus, lower emissivity and/or spectrally varying emissivity makes objects appear different temperatures than they are

How do we more accurately identify materials with a Thermal Imaging Camera?

By spectrally selective imaging to more accurately image emitted power.

Approach

Desired function of our filters



- Divide camera imaging range (7 μm -14 μm) into four equal energy/frequency bands
- Transmit one of four bands while reflecting the other three.
- Design material specific filters to only transmit where select materials are highly absorptive.

Distributed Bragg Reflector (DBR)

Ge (200nm)
AlO _x (400nm)
Ge (200nm)
AlO _x (400nm)
Ge (200nm)


- A DBR is a spectrally selective, highly reflective optical structure
- Each interface partially reflects incident light resulting in constructive interference
- Tunability of reflection band limited by material choice and layer thicknesses

Aperiodic Distributed Bragg Reflector (aDBR)

Ge (252 nm)
AlO _x (618 nm)
Ge (785 nm)
AlO _x (249 nm)
Ge (685 nm)
AlO _x (212 nm)
Ge (332 nm)
substrate

- Aperiodic DBR designs allow for different thicknesses of the various materials
- With different thicknesses and material arrangements, the aDBR can reflect, transmit, and absorb at different wavelengths
- Great tunability in design!
- Designs are too complex for intuition

Inverse Design Process




Inverse Design Algorithm


- Written in Python with API TensorFlow
- Target spectrum chosen to match desired response
- aDBR structure is randomly initialized
- Reflection spectra of aDBR is calculated via Transfer Matrix Method (design spectrum) and compared to target spectrum
- Based on difference b/w them (error) structure is modified
- Repeat progress until set iterations

Results


Simulated Structures



- Each narrow-band filter enables the thermal camera to detect its own peak emitted power
- Employing each of these filters with the thermal camera will allow for spectrally selective imaging
 - More accurate temperatures
 - More accurate material identification



Filter Application onto TIC



Current Status on Filter Growth



Image taken by Josh Nordlander

- Due to backorder on substrates, filter growth had been delayed.
- Currently during the filter's growth process, the surfaces start to become roughened/delaminated.
- Work is ongoing to reduce strain in the films.

References and Acknowledgements

- M. He, J.R. Nolen, J. Nordlander, A. Cleri, N.S. McIlwaine, Y. Tang, G. Lu, T.G. Folland, B.A. Landman, J.P. Maria, J.D. Caldwell, "Deterministic Inverse Design of Tamm Plasmon Thermal Emitters with Multi-Resonant Control", *Nature Materials*, in revision (2021).
- The authors would like to thank VINSE and the National Science Foundation (Grant #1852157) for their support.